

**Simultaneous Analysis of Cypermethrin and Fenvalerate  
Residues in Tomato after Field Treatment by Gas  
Chromatography**

**By**

**Mahgoub Ibrahim Shinger**

**B.Sc. in applied chemistry (IUA) 2003**

**A thesis submitted to the University of Khartoum  
for the Degree of M.Sc. in Analytical Chemistry**

**Supervisor**

**Dr. Abdalla Ahmed Elbashir**

**Co-supervisor**

**Dr. Hassan El Obeid Ahmed**

**Faculty of Science**

**University of Khartoum**

**July 2011**

## Table of contents

Contents		Page No
	Table of contents	I
	List of Tables	III
	List of Figures	IV
	List of abbreviation	V
	Acknowledgements	VI
	Arabic abstract	VII
	English abstract	VIII
Chapter one		
1	Introduction	1
1.1	History of pesticides	1
1.2	Pesticides health effects	2
1.2.1	Farmers and workers	2
1.2.2	Consumers	3
1.2.3	Public	3
1.3	Classification of pesticides	4
1.3.1	Classification of insecticides	4
1.3.1.1	Inorganic insecticides	5
1.3.1.2	Plant insecticides	5
1.3.1.3	Synthetic organic insecticides	6
1.3.1.3.1	Cypermethrin	7
1.3.1.3.1.1	Physical properties	8
1.3.1.3.1.2	Toxicological effects	8
1.3.1.3.1.3	Chemical structure of optical isomers	9
1.3.1.3.2	Fenvalerate	11
1.3.1.3.2.1	Physical properties	12
1.3.1.3.2.2	Toxicological effects	12
1.3.1.3.2.3	Chemical structure of optical isomers	13
1.4	Analytical techniques for determination of cypermethrin and fenvalerate residues in vegetables	15
1.4.1	Sample preparation and determination	15
1.4.1.1	Extraction and removal of co-extracted water	16
1.4.1.1.1	Organic solvent extraction	16
1.4.1.1.2	Matrix solid-phase dispersion extraction (MSPD)	19
1.4.1.1.3	Supercritical fluid extraction (SFE)	20
1.4.1.1.4	Pressurized fluid extraction (PFE)	20
1.4.1.1.5	Microwave-assisted extraction (MAE)	20

1.4.1.2	Sample cleanup	21
1.4.1.2.1	Liquid–liquid partitioning	21
1.4.1.2.2	Gel permeation chromatography (GPC)	21
1.4.1.2.3	Solid-phase extraction (SPE)	22
1.4.1.2.3.1	C <sub>18</sub> solid-phase extraction cleanup	22
1.4.1.2.3.2	Aminopropyl (NH <sub>2</sub> ), primary secondary amine (PSA) solid-phase extraction cleanup	22
1.4.1.2.3.3	Tandem GCB-NH <sub>2</sub> , GCB-PSA, and GCB-SAX-PSA SPE cleanup	23
1.4.1.2.3.4	Alumina, florisil, and silica solid-phase extraction cleanup	23
1.4.1.3	Analytical method for determination of cypermethrin and fenvalerate	24
1.5	Objectives	29
Chapter two		
2.	Experimental	30
2.1.	Apparatus	30
2.2.	Materials	30
2.2.1.	Solvents	30
2.2.2.	Saturated acetonitrile	30
2.2.3.	Deactivation of florisil	30
2.3.	Preparation of chromatographic column	30
2.4.	Preparation of calibration curves	31
2.5.	Sample preparation and fortification	31
2.6.	Extraction and cleanup procedures	31
2.7.	Analysis of the degradation of cypermethrin and fenvalerate in tomato	32
2.8.	Preparation of cypermethrin and fenvalerate formulations	32
Chapter three		
3.	Results and discussion	34
3.1.	Optimization of GC conditions	34
3.2.	Calibration curves	36
3.3.	Recoveries of spiked tomato	38
3.4.	Analysis of the degradation of cypermethrin and fenvalerate in tomato	39
3.5.	Analysis of cypermethrin and fenvalerate in formulations	43
4.	Conclusion	48
5.	References	49

## List Tables

	<b>Content</b>	<b>Page No</b>
Table 1	Identity of cypermethrin.	7
Table 2	Physical properties of cypermethrin.	8
Table 3	Identity of fenvalerate.	11
Table 4	Physical properties of fenvalerate.	12
Table 5	Properties of common solvents used in pesticide residue analysis at 20–25 °C	17
Table 6	Reported analytical procedures for recoveries of cypermethrin from fortified tomatoes.	25
Table 7	Reported analytical procedures for recoveries of fenvalerate from fortified tomatoes.	27
Table 8	The retention times of cypermethrin and fenvalerate.	35
Table 9	Calibration data for cypermethrin.	36
Table 10	Calibration data for fenvalerate.	37
Table 11	Analytical recoveries, LOD, LOQ and R <sup>2</sup> values of cypermethrin and fenvalerate in spiked tomato.	39
Table 12	Cypermethrin and fenvalerate residues at various time intervals following its application.	40
Table 13	In concentrations (mg/kg) of cypermethrin and fenvalerate on tomato after field treatment.	41
Table 14	Degradation rate constants of cypermethrin and fenvalerate on tomato after field treatment.	43
Table 15	Influence of different solvents used for cleanup on cypermethrin recoveries from formulation.	44
Table 16	Influence of different solvents used for cleanup on fenvalerate recoveries from formulation.	44

## List Figures

	<b>Content</b>	<b>Page No</b>
Figure 1	Chemical structures of cypermethrin isomers.	10
Figure 2	Chemical structures of fenvalerate isomers.	13
Figure 3	Schematic diagram explain the steps for determination of cypermethrin and fenvalerate residues in vegetables	14
Figure 4	Schematic diagram explain sequence of steps required for analysis of pesticides residues in vegetables	16
Figure 5	The chromatogram of standard stock solution of cypermethrin and fenvalerate analyzed by GC-ECD.	35
Figure 6	Calibration curve for cypermethrin.	36
Figure 7	Calibration curve for fenvalerate.	37
Figure 8	The chromatogram of blank tomato by GC-ECD.	38
Figure 9	Plot ln concentration (mg/kg) versus time for degradation of cypermethrin on tomato after field treatment.	42
Figure 10	Plot ln concentration (mg/kg) versus time for degradation of fenvalerate on tomato after field treatment.	42
Figure 11	Chromatogram of cypermethrin and fenvalerate detected on tomato after 1h from spraying.	45
Figure 12	Chromatogram of cypermethrin and fenvalerate detected on tomato after 7 days from spraying.	45
Figure 13	Chromatogram of cypermethrin and fenvalerate detected on tomato after 10 days from spraying.	45
Figure 14	Chromatogram of cypermethrin and fenvalerate detected on tomato after 13 days from spraying.	46
Figure 15	Chromatogram of cypermethrin and fenvalerate detected on tomato after 16 days from spraying.	46
Figure 16	Chromatogram of cypermethrin and fenvalerate detected on tomato after 19 days from spraying.	46
Fig 15	Chromatogram of cypermethrin and fenvalerate extracted from formulation using ethyl acetate: cyclohexane (1:1/v: v).	47
Fig 16	Chromatogram of cypermethrin and fenvalerate extracted from formulation using cyclohexane.	47
Fig 17	Chromatogram of cypermethrin and fenvalerate extracted from formulation using hexane.	47

## List of abbreviations

AD	Anno Domini
ASE	Accelerated Solvent Extraction
BC	Before Christ
Class I	Most Toxic
Class IV	Least Toxic
ECD	Electron Capture Detector
ELCD	Electrolytic Conductivity Detector
Envi-Carb	Environmental Carbon
EPA	Environmental Protection Agency
FDA	Food and Drug Administration
FPD	Flame Photometric Detector
FTD	Flame Thermionic Detector
GC	Gas Chromatography
GCB	Graphitized Carbon Black
GPC	Gel Permeation Chromatography
HPLC	High Performance Liquid Chromatography
IUPAC	International Union of Pure and Applied Chemists
LOD	Limit Of Detection
LOQ	Limit Of Quantification
MAE	Microwave Assisted Extraction
MDE	Macro Porous Diatomaceous Earth
MRLs	Maximum Residue Limits
MRM	Multiresidue Multiclass Method
MS	Mass Spectrometer
MSPD	Matrix Solid Phase Dispersion
NPD	Nitrogen Phosphorus Detector
PFE	Pressurized Fluid Extraction
PSA	Primary Secondary Amine
SAX	Strong Anion Exchange
SDVB	Styrene-Di Vinyl Benzene
SFE	Supercritical Fluid Extraction
SPE	Solid Phase Extraction
UV	Ultraviolet
WHO	World Health Organization

## ACKNOWLEDGEMENTS

- I thank Allah, Almighty, who helped me to complete this work.
- Special thanks to Dr. Abdalla Ahmed Elbashir and Dr. Hassan EL Obeid Ahmed, project supervisors.

This thesis would not have been possible unless their direction, guidance and encouragements.

- I wish to express my sincere gratitude and appreciation to Dr. Sheihk Idris Ahmed and Dr. AbdelRahman Hassan El Nadi.
- It is a pleasure to thank those who made this thesis possible, to Mr. Abuzar and other member off staff of the central lab of the chemistry department-Khartoum University.
- A thank go to International University of Africa, who permitted me the chance of education, and also the staff of the chemistry department.
- Thanks go to my family, who took part of my education honestly.
- Lastly, I offer my regards and blessing to all those who supported me in any respect.

## الخلاصة

في هذه الدراسة تم تقدير آني لمتبقيات السايبرمثرين و الفينفالريت في فاكهة الطماطم المزروعة في السودان في منطقة شرق الخرطوم بإستخدام جهاز كروماتوغرافيا الغاز المزود بمكشاف الإلكترونات.

الطريقة كانت خطية على مدى ٠,٠٧٥ - ٠,٠٠٩ ملجم/مللتر و ٠,٧٥ - ٠,٠٣٧ ملجم/مللتر لكل من السايبرمثرين و الفينفالريت على التوالي. وجد أنّ حدي الكشف و القياس الكمي هما ٠,٠٠٣ ، ٠,٠١ ملجم/مللتر و ٠,٠١٤ ، ٠,٠٤٧ ملجم/مللتر لكل من السايبرمثرين و الفينفالريت على التوالي.

كما وجد أنّ قيمتي الإسترجاع للسايبرمثرين و الفينفالريت من الطماطم المقواه هما ٩٧ %  $\pm$  ٧,٥ و ٩٩,٥ %  $\pm$  ١,٨ على التوالي.

أجريت دراسة حركية إضمحلال كل من المبيدين و أدى التقييم النهائي لبيانات الحركية الى وجود حركيات الرتبة الأولى بالنسبة لإضمحلالهما على فاكهة الطماطم. و من ثم قُدِّمَ تبرير هذه النتائج.

أُجري تحليل للتركيبية التجارية للمبيدين و أُختبرت مذيبيات مختلفة. فحصل على المستخلص الأكثر نقاوة و الأعلى قيم إسترجاع بإستخدام الهكسان الحلقي كمذيب تجزئة.



## Abstract

In this study simultaneous determination of cypermethrin and fenvalerate residues in tomato fruit (*Lycopersicon esculentum*) grown in The Sudan, in Khartoum area, was carried out using gas chromatography with electron capture detector (GC-ECD).

The method was linear in the range of 0.075 – 0.009 mg/ml and 0.75 – 0.037 mg/ml for cypermethrin and fenvalerate respectively. The limit of detection (LOD) and limit of quantification (LOQ) were found to be 0.003, 0.01 mg/ml and 0.014, 0.047 mg/ml for cypermethrin and fenvalerate respectively.

The recoveries of cypermethrin and fenvalerate spiked to tomato were 97%  $\pm$  7.5 and 99.5 %  $\pm$  1.8, respectively.

The kinetic study of the degradation of both pesticides was performed and the ultimate evaluation of the kinetic data revealed a first order kinetics with respect to their degradation in tomato fruit and an explanation was put forward to account for the results.

Formulation of cypermethrin and fenvalerate were analyzed, different solvents were tested. The cleanest extract and the best recoveries were found using cyclohexane as partitioning solvent.